U.S. DEPARTMENT OF COMMERCE
National Technical Information Service

PB-283 816

Winter Temperature Requirements for Maturation and Spawning of Yellow Perch 'Perca flavescens' (Mitchill)

Environmental Research Lab -Duluth, Minn

1977

TECHNICAL REPORT DATA
(Please read Instructions on the reverse before completing)

1. REPORT NO. EPA-600/J-77-126	2. JOURNAL ARTICLE
3. RECIPIENT'S ACCESSION NO. PB283816	
4. TITLE AND SUBTITLE: Winter Temperature Requirements for Maturation and Spawning of Yellow Perch Perca flavescens (Mitchill)	5. REPORT DATE
	6. PERFORMING ORGANIZATION CODE
7. AUTHOR(S): B. R. Jones, K. E. F. Hokanson, and J. H. McCormick	8. PERFORMING ORGANIZATION REPORT NO.
9. PERFORMING ORGANIZATION NAME AND ADDRESS: U.S. Environmental Protection Agency, Environmental Research Laboratory-Duluth, 6201 Congdon Boulevard, Duluth, MN 55804	10. PROGRAM ELEMENT NO. 1BA608
	11. CONTRACT/GRANT NO.
12. SPONSORING AGENCY NAME AND ADDRESS: Same	13. TYPE OF REPORT AND PERIOD COVERED
	14. SPONSORING AGENCY CODE EPA-600/03

15. SUPPLEMENTARY NOTES: IN: M. Marios (ed.), Towards a Plan of Actions for Mankind, Volume 3, Biological Balance and Thermal Modifications; Proceedings of the World Conference, Permagon Press, New York, N.Y. pp. 189-192.

16. ABSTRACT

Groups of sexually mature yellow perch were held during the winter at constant temperatures of 4 and 6 C for 123, 143, 164, 185, and 213 days, and of 8 and 10 C for 157, 201, and 241 days. The winter temperatures were raised to 20 C at a rate of 2 C per week at the end of each duration. The greatest reproductive success occurred in the group exposed to 4 C for 185 days at a median spawning temperature of 12 C. High spawning temperatures were associated with shorter durations and lower reproductive success. Reproductive success was lower at maturation temperatures above 4 C and at durations departing from 185 days.

17. KEY WORDS AND DOCUMENT ANALYSIS

a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
Temperature Fish Reproduction	Seasonal variations Reproductive success Spawning temperature	06-F

18. DISTRIBUTION STATEMENT: RELEASE TO PUBLIC	19. SECURITY CLASS *(This Report)* UNCLASSIFIED	21. NO. OF PAGES 5
	20. SECURITY CLASS *(This page)* UNCLASSIFIED	22. PRICE PC A02 MF A01

EPA Form 2220-1 (Rev. 4-77) PREVIOUS EDITIC

INSTRUCTIONS

1. **REPORT NUMBER**
   Insert the EPA report number as it appears on the cover of the publication.

2. **LEAVE BLANK**

3. **RECIPIENTS ACCESSION NUMBER**
   Reserved for use by each report recipient.

4. **TITLE AND SUBTITLE**
   Title should indicate clearly and briefly the subject coverage of the report, and be displayed prominently. Set subtitle, if used, in smaller type or otherwise subordinate it to main title. When a report is prepared in more than one volume, repeat the primary title, add volume number and include subtitle for the specific title.

5. **REPORT DATE**
   Each report shall carry a date indicating at least month and year. Indicate the basis on which it was selected *(e.g., date of issue, date of approval, date of preparation, etc.)*.

6. **PERFORMING ORGANIZATION CODE**
   Leave blank.

7. **AUTHOR(S)**
   Give name(s) in conventional order *(John R. Doe, J. Robert Doe, etc.)*. List author's affiliation if it differs from the performing organization.

8. **PERFORMING ORGANIZATION REPORT NUMBER**
   Insert if performing organization wishes to assign this number.

9. **PERFORMING ORGANIZATION NAME AND ADDRESS**
   Give name, street, city, state, and ZIP code. List no more than two levels of an organizational hirearchy.

10. **PROGRAM ELEMENT NUMBER**
    Use the program element number under which the report was prepared. Subordinate numbers may be included in parentheses.

11. **CONTRACT/GRANT NUMBER**
    Insert contract or grant number under which report was prepared.

12. **SPONSORING AGENCY NAME AND ADDRESS**
    Include ZIP code.

13. **TYPE OF REPORT AND PERIOD COVERED**
    Indicate interim final, etc., and if applicable, dates covered.

14. **SPONSORING AGENCY CODE**
    Insert appropriate code.

15. **SUPPLEMENTARY NOTES**
    Enter information not included elsewhere but useful, such as: Prepared in cooperation with, Translation of, Presented at conference of, To be published in, Supersedes, Supplements, etc.

16. **ABSTRACT**
    Include a brief *(200 words or less)* factual summary of the most significant information contained in the report. If the report contains a significant bibliography or literature survey, mention it here.

17. **KEY WORDS AND DOCUMENT ANALYSIS**
    (a) DESCRIPTORS - Select from the Thesaurus of Engineering and Scientific Terms the proper authorized terms that identify the major concept of the research and are sufficiently specific and precise to be used as index entries for cataloging.

    (b) IDENTIFIERS AND OPEN-ENDED TERMS - Use identifiers for project names, code names, equipment designators, etc. Use open-ended terms written in descriptor form for those subjects for which no descriptor exists.

    (c) COSATI FIELD GROUP - Field and group assignments are to be taken from the 1965 COSATI Subject Category List. Since the majority of documents are multidisciplinary in nature, the Primary Field/Group assignment(s) will be specific discipline, area of human endeavor, or type of physical object. The application(s) will be cross-referenced with secondary Field/Group assignments that will follow the primary posting(s).

18. **DISTRIBUTION STATEMENT**
    Denote releasability to the public or limitation for reasons other than security for example "Release Unlimited." Cite any availability to the public, with address and price.

19. & 20. **SECURITY CLASSIFICATION**
    DO NOT submit classified reports to the National Technical Information service.

21. **NUMBER OF PAGES**
    Insert the total number of pages, including this one and unnumbered pages, but exclude distribution list, if any.

22. **PRICE**
    Insert the price set by the National Technical Information Service or the Government Printing Office, if known.

PB 283 816

Reprinted from

Proceedings of the World Conference

TOWARDS A PLAN OF ACTIONS FOR MANKIND

VOLUME 3

BIOLOGICAL BALANCE AND THERMAL MODIFICATIONS

Edited by

M. MAROIS

PERGAMON PRESS . OXFORD and NEW YORK . 1977

EPA-600/J-77-126
JOURNAL ARTICLE

WINTER TEMPERATURE REQUIREMENTS FOR MATURATION AND SPAWNING OF YELLOW PERCH *PERCA FLAVESCENS* (MITCHILL)

B. R. JONES, K. E. F. HOKANSON, AND J. H. MCCORMICK

US Environmental Protection Agency, Environmental Research Laboratory–Duluth, Duluth, Minnesota, USA

INTRODUCTION

This is a report on a laboratory study of the winter temperature requirements for maturation and spawning of yellow perch, *Perca flavescens* (Mitchill).

The study was conducted to determine if addition of enough heat to the habitat of yellow perch to raise the winter temperature or shorten the period at usual low winter temperatures would be harmful to reproduction of this species. Information on this question would help predict the effects of heat addition to the aquatic environment and promote the adoption of water temperature standards related to seasonal biological needs.

Reports in the literature indicate that, for some fish, low temperatures in winter are essential to completion of gametogenesis. Examples are oogenesis in the four-spined stickleback, *Apeltes quadracus* (Merriman and Schedl, 1941) and spermatogenesis in the lake chub, *Couesius plumbeus* (Ashan, 1966).

Levels of reproductive success in relation to maturation temperatures have been reported for the fathead minnow, *Pimephales promelas* (Brungs, 1971) and the brook trout, *Salvelinus fontinalis* (Hokanson *et al.*, 1973) but not, to our knowledge, for the yellow perch.

Studies of gonads of yellow perch, however, show that in nature spermatogenesis starts in August and is completed between October and December, with the testes remaining full of spermatozoa until spawning (Turner, 1919; LeCren, 1951) and that ovaries increase in size from August to the time of spawning (LeCren, 1951).

This study determined the effects of various durations and temperatures on spawning and embryo survival in yellow perch.

METHODS

The study was conducted in 16 rectangular fiberglass tanks, 152 × 26 × 28 cm deep. The test temperatures were maintained by a chiller and by electronic controllers and valves that mixed hot and cold water in head boxes. The fish were held at the Duluth, Minnesota, photoperiod.

At the start of the test in October, 16 fish were placed in each tank. At the start of the spring increase in temperature, the number of fish was reduced and the sex ratio was adjusted to 1 male to 3 females or a maximum of 3 males and 7 females per tank. Previous experience with yellow perch in the laboratory indicated they would spawn in the rectangular fiberglass tanks without a special substrate.

The total number of eggs was obtained from the volume of spawn and the average number of eggs (70) per milliliter. The percent viable eggs was obtained by incubating 3

TABLE 1. Test temperatures and durations

Winter		Spring[1]	
Constant temperature (°C)	Duration from Oct. 30 (days)	Increase from winter[2] (°C)	Duration of increase (days)
4	123	18.6	43
	143	19.7	55
	164	17.8	42
	185	12.8	25
	213	7.3	8
6	123	—	—
	143	19.8	56
	164	16.0	35
	185	13.5	27
	213	7.7	6
8	157	18.6	30
	201	18.2	29
	241	7.9	—
10	157	19.0	33
	201	10.2	—
	241	10.2	—

[1] Increased at rate of 2°C per week
[2] Temperature at last spawning

subsamples of about 100 eggs each at 12 degrees centigrade. Egg viability was determined at formation of the neural keel.

The total temperature experience of the fish from October 30, to the last spawning in spring is summarized in Table 1. The fish were exposed over winter to constant temperatures of 4, 6, 8 and 10 degrees centigrade and several durations at these temperatures starting from October 30. They were exposed to 5 durations ranging from 123–213 days at 4 and 6 degrees and 3 durations ranging from 157–241 days at 8 and 10 degrees. At the end of the durations the temperature was increased at the rate of 2 degrees centigrade per week. By the time of last spawning, temperatures had risen to those shown in Table 1 as spring temperatures.

RESULTS

The results of the 16 combinations of temperature and duration on production of eggs are summarized in Table 2. The data are averaged for the shorter and longer durations at 4

and 6° and all durations at 8 and 10°. Some or all spawning occurred before the start of the spring increase in temperature at the longest durations at each temperature (213 days at 4 and 6°, 241 days at 8°, and 201 and 241 days at 10°). No spawning occurred at 6° and 123 days.

The average total numbers of eggs per female were greatest and about the same at the longer durations at 6° and the shorter and longer durations at 6°. Fewer eggs were produced at the shorter durations at 4° and fewer eggs were produced at the durations at 8 and 10°. Size of female may have had some influence on the total number of eggs produced (range in weight at termination was 42–200 g).

The average percent viable eggs per female, for the grouped durations, was highest (74%) at the longer durations at 4°, somewhat lower (41%) at the longer durations at 6°, and lower at the shorter durations at 4 and 6° and at 8 and 10°. The maximum percentages of viable eggs per female at any one combination of temperature and duration were 93% at

TABLE 2. Effects on Total Number of Eggs and Percent Viable Eggs per Female

°C	Duration (days)	Total eggs (×1000)	Viable eggs (%)
4	123, 143, 164	5	17
	185, 213	11	74
6	123, 143, 164	10	27
	185, 213	11	41
8	157, 201, 241	7	31
10	157, 201, 241	1	4

4° and 185 days' duration, 64% at 4° and 213 days' duration, and 64% at 6° and 164 days' duration.

Other responses which support these results were the number of spawnings per female and the quality of spawn. The number of spawnings per female were fewer and the quality of spawn was poorer at the shorter durations at 4 and 6° and at all durations at 8 and 10°. Good quality spawn consisted of complete, usually large (<100 m/l), accordian-folded ribbons as has been described for yellow perch (Mansueti, 1974). Poor quality spawn consisted of small volumes (10–100 ml) of fragments of ribbons.

The median spawning date at all test conditions ranged from May 4 to June 20 except one on April 12, and were within the spawning periods reported for yellow perch in natural waters (March to April, Mansueti, 1964; and May to June, Wells, 1970).

The average spawning temperatures at all test conditions ranged from 5.8 to 19°C. The greatest reproductive success, 93%, viable eggs at 4° and a duration of 185 days, occurred at an average spawning temperature of 11.3 degrees and within a range from 7.5–12°C. Reproductive success was poor when spawning occurred at temperatures above 15°.

The highest spawning temperatures occurred at the shortest durations at each temperature to suggest (since a temperature rise was not needed to trigger spawning) a need for higher "compensatory" spring temperatures to meet a total temperature requirement for maturation. Reproductive success was lowest under these conditions.

The conditions tested are related to percentage ranges of viable eggs in Table 3 to illustrate implications for fish production and in turn implications for water temperature

criteria. Only one winter temperature regimen (4° for 185 days) falls in the 75–100% range, two (4° for 213 days and 6° for 164 days) fall in the 50–75% range, and five fall in the

TABLE 3. Potential for Production

% Viable eggs			
100–75	75–50	50–25	25–0
°C/days	°C/days	°C/days	
4° 185	4° 213	4° 164	
	6° 164	6° 185	All
		6° 213	other
		8° 210	treatment
		8° 241	

25–50% range. The other 8 temperature regimens tested produced less than 25% viable eggs. It might be noted that durations of 213 days at 4 and 6° and 201 and 241 days at 8° are not likely to occur at the latitude at which the fish were collected and tested (47°N latitude).

CONCLUSION

It appears from this study that yellow perch need a low temperature in winter for maturation and spawning and that yellow perch suffer increased impairment of reproductive success as winter temperatures increase from a low of approximately 4°C and as durations of exposure to low temperatures decrease from approximately 185 days.

REFERENCES

Ahsan, Syed Nazar (1966) Effects of temperature and light on the cyclical changes in the spermatogenetic activity of the lake chub, *Couesius plumbeus* (Agassiz). *Can. J. Zool.* **44**, 161–171.

Brungs, W. A. (1971) Chronic effects of constant elevated temperature on the fathead minnow (*Pimephales promelas* Rafinesque). *Trans. Am. Fish. Soc.* **100**, 659–664.

Hokanson, K. E. F., McCormick, J. H., Jones, B. R., and Tucker, J. H. (1973) Thermal requirements for maturation, spawning, and embryo survival of the brook trout, *Salvelinus fontinalis* (Mitchill). *J. Fish. Res. Board Can.* **30**, 975–984.

LeCren, E. D. (1951) The length–weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *J. Anim. Ecol.* **20**, 201–219.

Mansueti, A. J. (1964) Early development of the yellow perch, *Perca flavescens*. *Chesapeake Sci.* **5**, 46–66.

Merriman, D., and Schedl, H. P., (1941) The effects of light and temperature on gametogenesis in the four-spined stickleback, *Apeltes quardracus* (Mitchill). *J. Exp. Biol.* **12**, 413–449.

Turner, C. L. (1919) The seasonal cycle in the spermary of the perch. *J. Morph.* **32**, 681–711.

Wells, L. (1970) In: Part IV. Effects of temperature fluctuations on Lake Michigan fish. Physical and ecological effects of waste heat on Lake Michigan. *Rept. Great Lakes Fish. Lab.*, Bur. Comm. Fish., US Dept. Int., Ann Arbor, Michigan. 101 pp.

4<